

Jan 8, 2021

New York State Department  
of Environmental Conservation  
Division of Materials Management  
Pesticide Product Registration

**FIFRA 2(ee) Recommendation  
For Management of Conifer Blights  
in Field Production, Nursery and Outdoor Landscapes**

Classified for  
"RESTRICTED USE"  
in New York State  
under 6NYCRR Part 326

Doc id: 571578

**Torque™ Fungicide  
EPA Reg. No. 1001-87**

This recommendation is made as permitted under FIFRA Section 2(ee)  
and has not been submitted to or approved by the USEPA

**DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL and RESIDENTIAL LANDSCAPES**

PLANTS	DISEASE	APPLICATION
<b>Conifers</b> <ul style="list-style-type: none"> <li>• Juniper[s]</li> <li>• Pine[s]</li> </ul>	<b>Phomopsis Blight</b> <i>Phomopsis juniperovora</i>	For Conifer Blights, apply at a rate of 8-10 fl. oz./100 gallons as a full coverage spray to the point of drip. Apply as needed, beginning when conditions are favorable for disease development.
	<b>Tip blight</b> <i>Sirococcus strobolinus</i>	
	<b>Diplodia Tip Blight</b> <i>Sphaeropsis sapinea</i>	For diplodia tip blight and tip blight, make 3 applications every 14 days prior to major period of infection. Application should be started at the initiation of new growth.  For juniper phomopsis blight, make first application as soon as junipers start to grow, and repeat the applications every 14-21 days during periods of active growth.  Can be use as a tank mix or rotation 3336, Spectro, or Chlorothalonil.

*Torque* and *Spectro* are registered trademarks of Cleary Chemical LLC.

*Spirato* is a registered trademark of Nufarm Americas Inc.

*Pageant* is a registered trademark of BASF Professional and Specialty Solution